

## ***Gender Affirming Interventions for Gender Dysphoria***

**Effective:** September 1, 2022

**Next Review:** September 2022

**Last Review:** April 2022

### **IMPORTANT REMINDER**

Medical Policies are developed to provide guidance for members and providers regarding coverage in accordance with contract terms. Benefit determinations are based in all cases on the applicable contract language. To the extent there may be any conflict between the Medical Policy and contract language, the contract language takes precedence.

PLEASE NOTE: Contracts exclude from coverage, among other things, services or procedures that are considered investigational or cosmetic. Providers may bill members for services or procedures that are considered investigational or cosmetic. Providers are encouraged to inform members before rendering such services that the members are likely to be financially responsible for the cost of these services.

### **DESCRIPTION**

This policy addresses interventions for gender dysphoria, a marked incongruence between one's experienced/expressed gender and assigned gender.

### **MEDICAL POLICY CRITERIA**

**Notes:**

- Member contracts for covered services vary. Member contract language takes precedence over medical policy.
- The Washington Gender Affirming Treatment Act (SSB 5313; <https://legiscan.com/WA/bill/SB5313/2021>) addresses coverage for gender affirming treatment for relevant member contracts.
- This policy does not address the following interventions:
  - Psychotherapy, which may be considered medically necessary for gender dysphoria; and
  - Medications such as hormonal therapy (see Cross References).

- I. Gender affirming interventions for gender dysphoria may be considered **medically necessary** when either of the following criteria is met:
  - A. For *member contracts subject to Washington's Gender Affirming Treatment Act* (SSB 5313), all of the following criteria are met (1. – 5.):
    1. At least 2 licensed mental health professionals have diagnosed gender dysphoria, and recommend the intervention (Note: only 1 mental health professional referral is required for breast/chest surgery); and
    2. Six continuous months of hormone therapy as appropriate to the patient's gender goals unless hormones are not clinically indicated for the individual (Notes: hormonal therapy is not required prior to breast/chest surgery); and
    3. At least 6 months of living in a role that is congruent with the patient's identity; and
    4. The request is for treatment(s) as prescribed by the treating provider because of, related to, or consistent with a person's gender expression or identity and is prescribed in accordance with accepted standards of care; and
    5. Either of the following is met:
      - a. Request is for genital surgery and patient has reached the age of majority (defined as age 18 in Washington state); or
      - b. Request is not for genital surgery.
  - B. For *all other member contracts*, both of the following criteria are met (1. – 2.):
    1. All of following general criteria are met (a. – d.):
      - a. Age at least 18 years (Note: age requirement will not be applied to breast/chest surgery with documented provider determination of medical necessity of earlier intervention); and
      - b. At least 2 licensed mental health professionals have diagnosed gender dysphoria, and recommend the intervention (Note: only 1 mental health professional referral is required for breast/chest surgery); and
      - c. Six continuous months of hormone therapy as appropriate to the patient's gender goals unless hormones are not clinically indicated for the individual (Note: hormonal therapy is not required prior to breast/chest surgery); and
      - d. At least 6 months of living in a role that is congruent with the patient's identity.
    2. One or more of the following criteria are met:
      - a. The request is for any of the following procedures:
        - i. Clitoroplasty
        - ii. Hysterectomy (Note: Hysterectomy is considered medically necessary without routine review and is not required to meet Criterion I.B.1.)
        - iii. Labiaplasty

- iv. Breast/chest surgery (i.e., breast augmentation, breast reduction, mastectomy, mastopexy, nipple/areola reconstruction/repositioning)
- v. Metoidioplasty
- vi. Orchiectomy
- vii. Penectomy
- viii. Penile prostheses implantation
- ix. Phallic reconstruction/Phalloplasty
- x. Salpingo-oophorectomy
- xi. Scrotoplasty
- xii. Testicular prostheses implantation
- xiii. Urethroplasty
- xiv. Vaginectomy
- xv. Vaginoplasty

- b. Clinical documentation is submitted expressly documenting that the intervention would improve otherwise documented significant gender dysphoria and the request is for one or more of the following procedures:
  - i. Hair removal
  - ii. Hair transplantation
  - iii. Endometrial ablation when **all** of the following criteria are met:
    - a.) Hysteroscopy, sonohysterography (SIS), or pelvic ultrasound has been performed and report is provided; and
    - b.) Endometrial sampling or dilation and curettage (D&C) has been performed or is planned according to any of the following:
      - i.) Endometrial sampling or D&C has been performed and report is provided. The histopathology report is provided showing absence of endometrial hyperplasia or uterine cancer; or
      - ii.) Endometrial sampling or D&C has been performed and report is provided. The histopathology report is provided, but inadequate tissue was obtained for diagnosis; or
      - iii.) Cervical stenosis precludes endometrial sampling, and D&C is planned concomitantly with ablation procedure.

- II. Gender affirming surgical interventions for gender dysphoria are considered **not medically necessary** for gender dysphoria when either of the following is met:
  - A. For member contracts subject to *Washington's Gender Affirming Treatment Act* (SSB 5313), when Criterion I.A. is not met; or
  - B. For member contracts **not** subject to *Washington's Gender Affirming Treatment Act* (SSB 5313), when any of the following is met:

1. Interventions listed in Criterion I.B.2 that do not meet the medical necessity criteria listed in Criterion I.B.1.; or
2. Interventions not listed in Criterion I.B.2. including, but not limited to abdominoplasty, blepharoplasty, brow lift, calf implants, cheek/malar implants, chin/nose implants, collagen injections, face-lift, facial bone reduction, forehead lift, lip reduction, liposuction, neck tightening, panniculectomy, pectoral implants, reduction thyroid chondroplasty, rhinoplasty, suction-assisted lipoplasty of the waist, voice modification surgery, and revision to a previous gender affirming surgery because of dissatisfaction with the appearance.
3. Reversal of gender affirming interventions.

*NOTE: A summary of the supporting rationale for the policy criteria is at the end of the policy.*

## LIST OF INFORMATION NEEDED FOR REVIEW

### REQUIRED DOCUMENTATION:

The information below **must** be submitted for review to determine whether policy criteria are met. If any of these items are not submitted, it could impact the review and decision outcome:

- History and Physical/Chart Notes
  - Documentation of therapy requested if applicable
  - Documentation of patient capacity to make decisions/consent to treatment
- For medical treatment or mastectomy:
  - Documentation that a licensed mental health professional has diagnosed gender dysphoria
  - Documentation of length of time living as desired gender
  - Documentation of length of time therapy occurred including licensure of therapist
  - For patients under the age of 18, documented provider determination of medical necessity of earlier intervention
- For all surgical treatments:
  - Documentation that at least 2 licensed mental health professionals have diagnosed gender dysphoria and recommend surgical treatment
- For all surgical treatments, excluding breast/chest surgery:
  - Documentation of hormonal therapy including length of time administered
  - Documented treatment plan including if planned procedures are reversals
- For procedures in Criteria I.B.2.b.:
  - Documentation that the intervention would improve otherwise documented significant gender dysphoria
- In addition to the above, for endometrial ablation:

- Endometrial histopathological report or documentation cervical stenosis precludes endometrial sampling and D&C is planned to be completed concomitantly with ablation procedure.
- Hysteroscopy, sonohysterography (SIS), or pelvic ultrasound report

## CROSS REFERENCES

1. [Endometrial Ablation](#), Surgery, Policy No. 01
2. [Cosmetic and Reconstructive Surgery](#), Surgery, Policy No. 12
3. [Reconstructive Breast Surgery/Mastopexy, and Management of Breast Implants](#), Surgery, Policy No. 40
4. [Reduction Mammoplasty](#), Surgery, Policy No. 60
5. [Autologous Fat Grafting to the Breast and Adipose-derived Stem Cells](#), Surgery, Policy No. 182
6. [Medication Policy Manual](#), Do a find (Ctrl+F) and enter drug name in the find bar to locate the appropriate policy.

## BACKGROUND

This policy supports applicable professional association statements,<sup>[1-5]</sup> and is also intended to support the Affordable Care Act (ACA) Section 1557 final implementing regulations published on May 18, 2016, and applicable state requirements<sup>[6]</sup>.

### **MEDICAL AND SURGICAL INTERVENTIONS FOR GENDER DYSPHORIA**

A clinical diagnosis of gender dysphoria is required prior to intervention for the disorder. Gender affirming interventions typically include psychotherapy, hormone therapy and in some cases surgical procedures. Psychotherapy followed by hormone therapy is often the first medical treatment sought, although not all transgender individuals on hormone therapy choose to undergo gender affirming surgery.<sup>[2]</sup>

#### **Gender Dysphoria**

Gender dysphoria is defined by the Diagnostic and Statistical Manual of Mental Disorders DSM-5 Diagnostic Criteria as follows:<sup>[7]</sup>

#### **Gender Dysphoria in Children 302.6 (F64.2)**

- A. A marked incongruence between one's experienced/expressed gender and assigned gender, of at least 6 months' duration, as manifested by at least six of the following (one of which must be Criterion 1):
  1. A strong desire to be of the other gender or an insistence that one is the other gender (or some alternative gender, different from one's assigned gender).
  2. In boys (assigned gender), a strong preference for cross-dressing or simulating female attire; or in girls (assigned gender), a strong preference for wearing only typical masculine clothing and a strong resistance to wearing of typical feminine clothing.
  3. A strong preference for cross-gender roles in make-believe play or fantasy play.
  4. A strong preference for toys, games, or activities stereotypically used or engaged in by the other gender.
  5. A strong preference for playmates of the other gender.
  6. In boys (assigned gender), a strong rejection of typically masculine toys, games and activities and a strong avoidance of rough-and-tumble play; or in

girls (assigned gender), a strong rejection of typically feminine toys, games and activities.

7. A strong dislike of one's sexual anatomy.
  8. A strong desire for the primary and/or secondary sex characteristics that match one's experienced gender.
- B. The condition is associated with clinically significant distress or impairment in social, school, or other important areas of functioning.

*Specify if:*

**With a disorder of sex development** (e.g., a congenital adrenogenital disorder such as 255.2 [E25.0] congenital adrenal hyperplasia or 259.0 [E34.50] androgen insensitivity syndrome).

**Coding note:** Code the disorder of sex development as well as gender dysphoria.

### **Gender Dysphoria in Adolescents and Adults 302.85 (F64.1)**

- A. A marked incongruence between one's experienced/expressed gender and assigned gender, of at least 6 months' duration, as manifested by at least two of the following:
1. A marked incongruence between one's experienced/expressed gender and primary and/or secondary sex characteristics (or in young adolescents, the anticipated secondary sex characteristics).
  2. A strong desire to be rid of one's primary and/or secondary sex characteristics because of a marked incongruence with one's experienced/expressed gender (or in young adolescents, a desire to prevent the development of the anticipated secondary sex characteristics).
  3. A strong desire for the primary and /or secondary sex characteristics of the other gender.
  4. A strong desire to be of the other gender (or some alternative gender different from one's assigned gender).
  5. A strong desire to be treated as the other gender (or some alternative gender different from one's assigned gender).
  6. A strong conviction that one has the typical feelings and reactions of the other gender (or some alternative gender different from one's assigned gender).
- B. The condition is associated with clinically significant distress or impairment in social, occupational, or other important areas of functioning.

*Specify if:*

**With a disorder of sex development** (e.g., a congenital adrenogenital disorder such as 255.2 [E25.0] congenital adrenal hyperplasia or 259.0 [E34.50] androgen insensitivity syndrome).

**Coding note:** Code the disorder of sex development as well as gender dysphoria.

*Specify if:*

**Post transition:** The individual has transitioned to full-time living in the desired gender (with or without legalization of gender change) and has undergone (or is preparing to have) at least one cross-sex medical procedure or treatment regimen- namely regular cross-sex hormone treatment or gender reassignment

surgery confirming the desired gender (e.g., penectomy, vaginoplasty in the natal male; mastectomy or phalloplasty in the natal female).

## **Psychotherapy**

Psychotherapy provided by a mental health professional typically includes an initial assessment of gender identity and dysphoria, the historical development of gender dysphoric feelings, and severity of resulting stress caused by the condition.<sup>[4]</sup> The goal of therapy is to assess, diagnose, and discuss treatment options, if needed, and is typically required prior to hormone therapy and/or surgical treatment.

## **Hormone Therapy**

Hormone therapy is undertaken in order to feminize or masculinize individuals' bodies to conform to their desired gender identities. For transgender individuals, hormone replacement therapy (HRT) causes the development of many of the secondary sexual characteristics of their gender identity. Prescribed hormones differ depending upon the natal gender of the individual. For individuals seeking to feminize, hormone treatment may include estradiol, finasteride, and spironolactone. For individuals seeking to masculinize, hormone treatment may include androgenic hormones such as testosterone.

## **Surgical Interventions**

Surgical intervention for gender dysphoria differs depending upon the gender assigned at birth. For individuals who are feminizing, surgery may involve removal of the testicles and penis and the creation of a pseudo vagina, clitoris, and labia. Complications of feminizing genital surgery may include necrosis of the vagina and labia, neovaginal prolapse, fistulas from the bladder or bowel into the vagina, stenosis of the urethra, and small or short vaginas.<sup>[4, 8]</sup>

For individuals who are masculinizing, surgery may involve removal of the uterus, ovaries, and vagina, and creation of a neophallus and scrotum with scrotal and/or penile prostheses. The creation of a neophallus for these patients is a multistage reconstructive procedure. Currently, techniques for penile reconstruction procedures vary and complications may include frequent urinary tract stenoses and fistulas, donor site scarring and necrosis of the neophallus.<sup>[4, 9]</sup> In addition, breast size does not significantly decrease with hormonal therapy and as a result, masculinizing patients may choose to undergo mastectomy to remove breast tissue. For many patients this may be the only surgery undertaken.<sup>[4]</sup> Mastectomy may involve a complete resection of all breast tissue; however, the nipple/areola sparing technique is typically performed to preserve the nipple/areola. For those who are taking androgen hormones, menstruation usually ceases with the medication intervention alone. In those who experience continued uterine bleeding other hormonal regimes may be attempted, or endometrial ablation.<sup>[10]</sup>

There are various additional surgical procedures which may be sought in order to complete the physical gender transformation and align an individual to their gender identity. However, conflicting opinions exist regarding whether these procedures are essential in treating gender dysphoria.

The WPATH recommends that patients, “engage in 12 continuous months of living in a gender role that is congruent with their gender identity...” prior to gender reassignment surgery so that patients may socially adjust to their desired gender role.<sup>[4]</sup> WPATH notes that changing a

gender role may have personal and social consequences which should be adequately explored prior to undergoing an irreversible surgery.

## EVIDENCE SUMMARY

Evidence regarding interventions for gender dysphoria in transgender individuals primarily consists of systematic reviews consisting of small cohort studies. Randomized clinical trials (RCTs) comparing gender dysphoria interventions with no intervention are ideal. However, there are challenges in conducting RCTs for these interventions due to several factors, such as small patient populations, and ethical concerns regarding the high morbidity and mortality rates associated with no intervention. Therefore, large RCTs are not anticipated. This policy relies on the following systematic reviews and non-randomized studies, as well as professional association recommendations to support applicable federal and state requirements.

### SYSTEMATIC REVIEWS

Wernick (2019) published a systematic review of the psychological benefits of gender affirming surgery. A total of 33 studies met inclusion criteria. The key concepts searched were quality of life, gender-confirmation surgical procedures, and transgender persons. Sixteen of the identified studies addressed compared pre- and post-surgical data, while 17 studies compared between-group differences. No meta-analysis was completed. Most studies demonstrated a trend of better mental health in transgender individuals who underwent surgeries, but not all reported improvements were statistically significant. The systematic review concluded that gender affirming surgery may lead to psychological benefits for individuals with gender dysphoria and that more research is needed to understand the factors that contribute to the outcomes following these surgeries.

Berli (2017) published a review of the available literature regarding facial gender confirmation surgery (FGCS).<sup>[11]</sup> The literature search went through December, 2016. The evidence was evaluated using the Oxford Centre for Evidence-Based Medicine suggestions for levels of evidence. Based on their findings, Berli and colleagues recommended that the next World Professional Association for Transgender Health (WPATH) Standards of Care version should include specific FGCS procedures. The authors also recommended replacing the historical term, facial feminization surgery (FFS) with more inclusive terminology – facial gender confirmation surgery. The body of evidence regarding FGCS is limited to case reports and case series. The authors found most data did not include quality-of-life outcome measures, and when reported, standardized instruments were not utilized. FGCS procedures were categorized by the authors as structural (e.g., forehead reconstruction, rhinoplasty), and secondary nonstructural procedures (e.g., blepharoplasty, upper lip shortening techniques). The review was limited by the paucity of data on FGCS as a treatment for gender dysphoria. In addition, methodological limitations of the review included but were not limited to, lack of transparent study selection and a transparent, comprehensive assessment of study quality and risk of bias. These limitations prohibit conclusions about overall health outcomes.

In 2009, Murad assessed quality of life and other psychosocial outcomes of transgendered individuals with gender identity disorder (GID), receiving hormonal therapy as part of gender affirmation surgery.<sup>[12]</sup> Twenty-eight cohort studies were included in the review which included pooled data from 1,833 patients with GID (1,093 male-to-female [MTF] and 801 female-to-male [FTM]). Significant improvements were reported after gender affirmation compared to pre-treatment status: 80% of patients reported improvement in gender dysphoria (95% CI = 68-89%; 8 studies) 78% reported significant improvement in psychological symptoms (95% CI =

56-94%; 7 studies) 80% reported significant improvement in quality of life (95% CI = 72-88%; 16 studies); and 72% reported significant improvement in sexual function (95% CI = 60-81%; 15 studies). Significant study heterogeneity was reported for all outcomes. Although the authors acknowledge the low quality of evidence used in the analysis, gender affirmation that included hormonal interventions in patient with GID was thought to likely improve symptoms of gender dysphoria and overall quality of life.

In 2009, Elamin evaluated the use of sex steroids on cardiovascular risk in transgender individuals.<sup>[13]</sup> A total of 16 studies were included in the review with a total of 1,471 male-to-female (MTF) patients and 651 female-to-male (FTM) patients. Steroid use was associated with increased serum triglycerides in both MTF and FTM patients and a nonsignificant effect on HDL-cholesterol and systolic blood pressure in FTM patients. Authors noted that the quality of evidence was low due to methodological limitations of included studies, including but not limited to, heterogeneity of patient population and variable follow-up periods and uncontrolled study design.

### **Nonrandomized Studies**

Primary evidence is limited to cohort studies with a variety of methodological limitations, including but not limited to small sample size, short-term follow-up, lack of comparison group, and varied treatment methods. Despite these limitations, significant improvements in quality of life, psychological comorbidities, and sexual functioning were consistently reported in patients who received gender-confirming medical treatments.<sup>[14]</sup> Below are summaries of representative publications.

Morrison (2020) performed a prospective, international, multicenter cohort study to assess outcomes of facial feminization surgery.<sup>[15]</sup> Outcomes reported were facial feminization outcome scores, satisfaction, and cephalometric analysis of femininity. A total of 66 consecutive patients at two clinics were enrolled. The increase in median facial feminization score from pre- to six months post-surgery was statistically significant, from 47.2 to 80.6 ( $p < 0.0001$ ). Cephalometric measures, including glabellar angle, nasolabial angle, and forehead inclination, were significantly more feminine after surgery. Mean satisfaction, measured on a five-point Likert scale, was 3.0 at <1-month post-surgery and six months post-surgery. No general measures of quality of life or mental health were reported.

Imbimbo (2009) evaluated the clinical and psychosocial profile of male-to-female transgendered individuals who had undergone reconstructive surgery.<sup>[16]</sup> The average age of patients was 31 years old, 72% had high educational levels, half of patients' contemplated suicide at some point prior to surgery and 4% had attempted suicide. Improved sex life satisfaction was reported in 75% of patients, with almost all patients' reporting satisfaction with their new sexual status. Additional studies sought to evaluate the sociodemographic profile of transgender individuals with GID in an effort to better characterize and provide treatment for this population.<sup>[17]</sup>

Heylens (2014) assessed comorbidities and psychosocial factors at various phases of the gender affirmation process in 57 patients with GID.<sup>[18]</sup> The Symptom Checklist-90 (SCL-90) was administered at three time points: baseline, after the start of hormone therapy, and after sex reassignment surgery (SRS) (also known as [aka] gender affirmation surgery). Psychopathological parameters include overall psychoneurotic distress, anxiety, agoraphobia, depression, somatization, paranoid ideation/psychoticism, interpersonal sensitivity, hostility, and sleeping problems and the psychosocial parameters consist of relationship, living

situation, employment, sexual contacts, social contacts, substance abuse, and suicide attempt. The greatest improvement in psychoneurotic distress was observed after the initiation of hormone therapy ( $p < 0.001$ ). In addition, significant decreases in anxiety, depression, interpersonal sensitivity and hostility were reported after hormone therapy. No significant differences were observed in pre- and postoperative assessments.

Fisher (2013) described clinical and sociodemographic features of 140 transmen ( $n=48$ ) and transwomen ( $n=92$ ) with GID and without affirmation surgery.<sup>[19]</sup> The following assessment tests were administered: the Body Uneasiness Test (a self-rating scale exploring different areas of body-related psychopathology), Symptom Checklist-90 Revised (a self-rating scale to measure psychological state), and the Bem Sex Role Inventory (a self-rating scale to evaluate gender role). Authors reported that transmen displayed significantly better social functioning than transwoman.

Gorin-Lazard (2013) reported a case series which assessed a variety of gender dysphoria symptoms with hormonal treatment preceding gender affirmation surgery. Pre- and post-hormone treatment self-esteem (Social Self-Esteem Inventory), mood (Beck Depression Inventory), QoL (Subjective Quality of Life Analysis), and global functioning (Global Assessment of Functioning) scores were compared in 49 patients.<sup>[20]</sup> Hormone therapy was reported to be an independent factor in greater self-esteem, a reduction in depression, and improved QoL scores.

Gomez-Gil (2012) evaluated symptoms of social distress, anxiety and depression in 187 transgendered individuals.<sup>[21]</sup> Of those included in the study, 120 had undergone hormonal sex-reassignment (SR) (aka gender affirmation) treatment and 67 had not. Social anxiety was assessed with the Social Anxiety and Distress Scale (SADS) and depression and anxiety were assessed with the Hospital Anxiety and Depression Scale (HADS). The non-hormone group was reported to be significantly younger than the treatment group (mean age 25.9 vs. 33.6 years,  $p=0.001$ ) and was less likely to have undergone surgical interventions ( $p < 0.001$ ). After adjusting for confounding factors, the authors reported that patients who were receiving hormone treatment had significantly lower prevalence of depression, anxiety, and social anxiety than those not receiving hormones.

Johansson (2010) reported long-term (five-year) outcomes of transgendered individuals ( $n=42$ ) with GID who had completely transitioned ( $n=32$ ), were in progress ( $n=5$ ) or who were on hormone therapy ( $n=5$ ).<sup>[22]</sup> Authors reported that no patient regretted affirmation and clinicians rated the global outcome as favorable in 62% of the cases, compared to 95% according to the patients themselves, with no differences between the subgroups. At follow-up, more than 90% of patients reported stable or improved work situations, partner relations and sex-life. However, 5-15% of patients reported dissatisfaction with hormonal treatment, results of surgery, total gender affirmation procedure, or their present general health.

Asscheman (2011) evaluated the long-term (one-year) effects of cross-sex hormones in 966 male-to-female (MTF) and 365 female-to-male (FTM) transgendered individuals.<sup>[23]</sup> MTF patients received different doses of estrogen and cyproterone acetate and FTM patients received parenteral/oral testosterone esters or testosterone gel. Hormone treatment levels varied at pre-and post-surgical affirmation time points. High mortality rates were reported in the MTF group when compared to the general population (51%); however, this increased rate was due to non-hormone-related causes such as suicide, acquired immunodeficiency syndrome

(AIDS), cardiovascular disease, drug abuse and other unknown causes. No significant increase in mortality was observed in FTM patients compared to the general population.

## Summary

The evidence is limited by a lack of well-designed studies comparing the safety and effectiveness gender affirming surgery to no treatment or to hormone therapy alone. There are challenges in conducting these large studies, and therefore such studies are not expected in the near future. Although additional research is needed, the research addressing genital and chest surgeries has consistently suggested significant improvement in symptoms and overall quality of life. With regard to other surgeries, such as facial feminization and body contouring, evidence is insufficient to show an improvement in health outcomes.

## PRACTICE GUIDELINE SUMMARY

### WORLD PROFESSIONAL ASSOCIATION FOR TRANSGENDER HEALTH

The World Professional Association for Transgender Health (WPATH) is a multidisciplinary professional society representing the specialties of medicine, psychology, social sciences and law that has published clinical guidelines regarding health services for patients with gender disorders. In 2011, WPATH approved the update of their evidence and consensus-based guideline regarding, the *Standards of Care (SOC) for the Health of Transsexual, Transgender, and Gender Nonconforming Peoples, 7<sup>th</sup> Version*.<sup>[4]</sup> The 8<sup>th</sup> version is anticipated in 2021. WPATH guidelines describe surgical procedures as “irreversible changes to the body.” Therefore, WPATH guidelines recommend the appropriate care should be taken to ensure patients have sufficient time (at least 24 hours) to consider all the information and can provide informed consent. WPATH notes, “(t)hese surgeries may be performed once there is written documentation that this assessment has occurred and that the person has met the criteria for a specific surgical treatment. By following this procedure, mental health professionals, surgeons, and patients share responsibility for the decision to make irreversible changes to the body.”

### Physical Interventions for Adolescents

WPATH guidelines state that physical interventions for adolescents fall into three categories or stages:

1. Fully reversible interventions. These involve the use of GnRH analogues to suppress estrogen or testosterone production and consequently delay the physical changes of puberty. Alternative treatment options include progestins (most commonly medroxyprogesterone) or other medications (such as spironolactone) that decrease the effects of androgens secreted by the testicles of adolescents who are not receiving GnRH analogues. Continuous oral contraceptives (or depot medroxyprogesterone) may be used to suppress menses.
2. Partially reversible interventions. These include hormone therapy to masculinize or feminize the body. Some hormone-induced changes may need reconstructive surgery to reverse the effect (e.g., gynaecomastia caused by estrogens), while other changes are not reversible (e.g., deepening of the voice caused by testosterone).
3. Irreversible interventions. Reversible and irreversible interventions are outlined in the standards of care, specifying intervention sequencing in adolescents. It is also stated that “[t]wo goals justify intervention with puberty suppressing hormones: (i) their use gives adolescents more time to explore their gender nonconformity and other developmental issues;

and (ii) their use may facilitate transition by preventing the development of sex characteristics that are difficult or impossible to reverse if adolescents continue on to pursue sex reassignment.”

## **Referral for Surgery**

WPATH guidelines indicate that surgical interventions can be initiated by a referral from a qualified mental health professional. One or two referrals may be required depending upon the type of surgery requested. “The mental health professional provides documentation—in the chart and/or referral letter—of the patient’s personal and treatment history, progress, and eligibility.” WPATH guidelines specifically recommend the following:

- One referral from a qualified mental health professional is needed for breast/chest surgery (e.g., mastectomy, chest reconstruction, or augmentation mammoplasty).
- Two referrals—from qualified mental health professionals who have independently assessed the patient—are needed for genital surgery (i.e., hysterectomy/salpingo-oophorectomy, orchiectomy, genital reconstructive surgeries).

### **Criteria for Breast/Chest Surgery (One Referral)**

WPATH lists the following criteria for mastectomy and creation of a male chest in FTM patients:

1. Persistent, well-documented gender dysphoria;
2. Capacity to make a fully informed decision and to consent for treatment;
3. Age of majority in a given country;
4. If significant medical or mental health concerns are present, they must be reasonably well controlled.

Hormone therapy is not a prerequisite.

### **Criteria for Genital Surgery (Two Referrals)**

WPATH lists the following criteria for genital surgery:

1. Persistent, well-documented gender dysphoria;
2. Capacity to make a fully informed decision and to consent for treatment;
3. Age of majority in a given country;
4. If significant medical or mental health concerns are present, they must be reasonably well controlled.
5. 12 continuous months of hormone therapy as appropriate to the patient’s gender goals (unless hormones are not clinically indicated for the individual).

In addition, WPATH made specific recommendations regarding breast augmentation procedures:

### **Breast Augmentation**

The WPATH guidelines recommend patients undergo hormone therapy for a minimum of 12 months prior to augmentation surgery and lists specific criteria for breast augmentation (implants/lipofilling).

### **Other Procedures**

The WPATH guidelines state that “while most professionals agree that genital surgery and mastectomy cannot be considered purely cosmetic, opinions diverge as to what degree other surgical procedures (e.g. breast augmentation, facial feminization surgery) can be considered purely reconstructive.” The guidelines go on to say that “[a]lthough it may be much easier to see a phalloplasty or a vaginoplasty as an intervention to end lifelong suffering, for certain patients an intervention like a reduction rhinoplasty can have a radical and permanent effect on their quality of life, and therefore is much more medically necessary than for somebody without gender dysphoria.”

## Physical Effects of Hormone Therapy

The WPATH guidelines outline the time course of the physical changes that are induced by feminizing/masculinizing hormone therapy. Some of the effects of masculinizing hormones that are relevant to other interventions discussed here are:

- facial/body hair growth - expected onset three to six months, expected maximum effect three to five years
- cessation of menses - expected onset two to six months
- vaginal atrophy - expected onset three to six months, expected maximum effect one to two years
- deepened voice - expected onset 3 – 12 months, expected maximum effect one to two years

Some of the effects of feminizing hormones that are relevant to other interventions discussed here are:

- breast growth - expected onset three to six months, expected maximum effect two to three years
- decreased testicular volume - expected onset three to six months, expected maximum effect two to three years
- thinning and slowed growth of body and facial hair - expected onset 6 – 12 months, expected maximum effect over three years

## THE ENDOCRINE SOCIETY

In 2017, the Endocrine Society in conjunction with American Association of Clinical Endocrinologists, American Society of Andrology, European Society for Pediatric Endocrinology, European Society of Endocrinology, Pediatric Endocrine Society, and World Professional Association for Transgender Health published updated guidelines for the treatment of gender-dysphoric/gender-incongruent persons.<sup>[10]</sup> The guideline employed transparent methods for evidence review and for rating the quality of evidence. Guidelines were referenced as *recommendations* or *suggestions*, by the numbers 1 and 2, respectively. Evidence was ranked as very low-quality |⊕○○○; low quality |⊕⊕○○; moderate quality |⊕⊕⊕○; and high quality |⊕⊕⊕⊕. The consortium made the following statements:

### 1.0 Evaluation of Youth and Adults

1.1 We advise that only trained mental health professionals (MHPs) who meet the following criteria should diagnose gender dysphoria (GD)/ gender incongruence in adults: (1) competence in using the Diagnostic and Statistical Manual of Mental Disorders (DSM) and/or the International Statistical Classification of Diseases and Related Health

Problems (ICD) for diagnostic purposes, (2) the ability to diagnose GD/ gender incongruence and make a distinction between GD/gender incongruence and conditions that have similar features (e.g., body dysmorphic disorder), (3) training in diagnosing psychiatric conditions, (4) the ability to undertake or refer for appropriate treatment, (5) the ability to psychosocially assess the person's understanding, mental health, and social conditions that can impact gender-affirming hormone therapy, and (6) a practice of regularly attending relevant professional meetings. (Ungraded Good Practice Statement)

- 1.2. We advise that only MHPs who meet the following criteria should diagnose GD/gender incongruence in children and adolescents: (1) training in child and adolescent developmental psychology and psychopathology, (2) competence in using the DSM and/or the ICD for diagnostic purposes, (3) the ability to make a distinction between GD/gender incongruence and conditions that have similar features (e.g., body dysmorphic disorder), (4) training in diagnosing psychiatric conditions, (5) the ability to undertake or refer for appropriate treatment, (6) the ability to psychosocially assess the person's understanding and social conditions that can impact gender-affirming hormone therapy, (7) a practice of regularly attending relevant professional meetings, and (8) knowledge of the criteria for puberty blocking and gender-affirming hormone treatment in adolescents. (Ungraded Good Practice Statement)
- 1.3. We advise that decisions regarding the social transition of prepubertal youths with GD/gender incongruence are made with the assistance of an MHP or another experienced professional. (Ungraded Good Practice Statement).
- 1.4. We recommend against puberty blocking and gender-affirming hormone treatment in prepubertal children with GD/gender incongruence. (1 | ⊕ ⊕ ○ ○)
- 1.5. We recommend that clinicians inform and counsel all individuals seeking gender-affirming medical treatment regarding options for fertility preservation prior to initiating puberty suppression in adolescents and prior to treating with hormonal therapy of the affirmed gender in both adolescents and adults. (1 | ⊕ ⊕ ⊕ ○)

## 2.0 Treatment of Adolescents

- 2.1. We suggest that adolescents who meet diagnostic criteria for GD/gender incongruence, fulfill criteria for treatment, and are requesting treatment should initially undergo treatment to suppress pubertal development. (2 | ⊕ ⊕ ○ ○)
- 2.2. We suggest that clinicians begin pubertal hormone suppression after girls and boys first exhibit physical changes of puberty. (2 | ⊕ ⊕ ○ ○)
- 2.3. We recommend that, where indicated, GnRH analogues are used to suppress pubertal hormones. (1 | ⊕ ⊕ ○ ○)
- 2.4. In adolescents who request sex hormone treatment (given this is a partly irreversible treatment), we recommend initiating treatment using a gradually increasing dose schedule after a multidisciplinary team of medical and MHPs has confirmed the persistence of GD/gender incongruence and sufficient mental capacity to give informed consent, which most adolescents have by age 16 years. (1 | ⊕ ⊕ ○ ○).
- 2.5. We recognize that there may be compelling reasons to initiate sex hormone treatment prior to the age of 16 years in some adolescents with GD/ gender incongruence, even though there are minimal published studies of gender-affirming hormone treatments administered before age 13.5 to 14 years. As with the care of adolescents 16 years of age, we recommend that an expert multidisciplinary team of medical and MHPs manage this treatment. (1 | ⊕ ○ ○ ○)

2.6. We suggest monitoring clinical pubertal development every 3 to 6 months and laboratory parameters every 6 to 12 months during sex hormone treatment. (2 |⊕⊕○○)

### 3.0 Hormonal Therapy for Transgender Adults

3.1. We recommend that clinicians confirm the diagnostic criteria of GD/gender incongruence and the criteria for the endocrine phase of gender transition before beginning treatment. (1 |⊕⊕⊕○)

3.2. We recommend that clinicians evaluate and address medical conditions that can be exacerbated by hormone depletion and treatment with sex hormones of the affirmed gender before beginning treatment. (1 |⊕⊕⊕○)

3.3. We suggest that clinicians measure hormone levels during treatment to ensure that endogenous sex steroids are suppressed and administered sex steroids are maintained in the normal physiologic range for the affirmed gender. (2 |⊕⊕○○)

**3.4. We suggest that endocrinologists provide education to transgender individuals undergoing treatment about the onset and time course of physical changes induced by sex hormone treatment. (2 |⊕○○○)**

### 4.0 Adverse Outcome Prevention and Long-term Care

4.1. We suggest regular clinical evaluation for physical changes and potential adverse changes in response to sex steroid hormones and laboratory monitoring of sex steroid hormone levels every 3 months during the first year of hormone therapy for transgender males and females and then once or twice yearly. (2 |⊕⊕○○)

4.2. We suggest periodically monitoring prolactin levels in transgender females treated with estrogens. (2 |⊕⊕○○)

4.3. We suggest that clinicians evaluate transgender persons treated with hormones for cardiovascular risk factors using fasting lipid profiles, diabetes screening, and/or other diagnostic tools. (2 |⊕⊕○○)

4.4. We recommend that clinicians obtain bone mineral density (BMD) measurements when risk factors for osteoporosis exist, specifically in those who stop sex hormone therapy after gonadectomy. (1 |⊕⊕○○)

4.5. We suggest that transgender females with no known increased risk of breast cancer follow breast-screening guidelines recommended for non-transgender females. (2 |⊕⊕○○)

4.6. We suggest that transgender females treated with estrogens follow individualized screening according to personal risk for prostatic disease and prostate cancer. (2 |⊕○○○)

**4.7. We advise that clinicians determine the medical necessity of including a total hysterectomy and oophorectomy as part of gender-affirming surgery. (Ungraded Good Practice Statement)**

### 5.0 Surgery for Sex Reassignment and Gender Confirmation

5.1. We recommend that a patient pursue genital gender-affirming surgery only after the MHP and the clinician responsible for endocrine transition therapy both agree that surgery is medically necessary and would benefit the patient's overall health and/or well-being. (1 |⊕⊕○○)

5.2. We advise that clinicians approve genital gender affirming surgery only after completion of at least 1 year of consistent and compliant hormone treatment, unless hormone therapy is not desired or medically contraindicated. (Ungraded Good Practice Statement)

- 5.3. We advise that the clinician responsible for endocrine treatment and the primary care provider ensure appropriate medical clearance of transgender individuals for genital gender-affirming surgery and collaborate with the surgeon regarding hormone use during and after surgery. (Ungraded Good Practice Statement)
- 5.4. We recommend that clinicians refer hormone treated transgender individuals for genital surgery when: (1) the individual has had a satisfactory social role change, (2) the individual is satisfied about the hormonal effects, and (3) the individual desires definitive surgical changes. (1 | ⊕○○○)
- 5.5. We suggest that clinicians delay gender-affirming genital surgery involving gonadectomy and/or hysterectomy until the patient is at least 18 years old or legal age of majority in his or her country. (2 | ⊕⊕○○)
- 5.6. We suggest that clinicians determine the timing of breast surgery for transgender males based upon the physical and mental health status of the individual. There is insufficient evidence to recommend a specific age requirement. (2 | ⊕○○○)

## **AMERICAN COLLEGE OF OBSTETRICIANS AND GYNECOLOGY**

In 2017 and 2011, the American College of Obstetricians and Gynecology (ACOG) published committee opinions regarding care for transgender adolescents, and health care services for transgendered individuals, respectively.<sup>[24, 25]</sup> Although these guidelines are not based on evidence, ACOG does make the following statements:

“Obstetrician–gynecologists should be prepared to assist or refer transgender individuals for routine treatment and screening as well as hormonal and surgical therapies. Hormonal and surgical therapies for transgender patients may be requested, but should be managed in consultation with health care providers with expertise in specialized care and treatment of transgender patients.”

Regarding adolescents, ACOG highlights age-specific concerns with a focus on medical management.

“Consensus guidelines support initiating medical therapy after an adolescent has an established diagnosis of transgender identity and has reached Tanner stage II development.”

In addition, ACOG guidelines made specific recommendations regarding hormone therapy, surgery and screening for both female-to-male and male-to-female patients:

### **Female-to-Male Transgender Individuals**

#### Hormones

Methyltestosterone injections every 2 weeks are usually sufficient to suppress menses and induce masculine secondary sex characteristics. Before receiving androgen therapy, patients should be screened for medical contraindications and have periodic laboratory testing, including hemoglobin and hematocrit to evaluate for polycythemia, liver function tests, and serum testosterone level assessments (goal is a mid-normal male range of 500 microgram/dL), while receiving the treatment.

#### Surgery

Hysterectomy, with or without salpingo-oophorectomy, is commonly part of the surgical process. An obstetrician–gynecologist who has no specialized expertise in transgender

care may be asked to perform this surgery, and also may be consulted for routine reasons such as dysfunctional bleeding or pelvic pain. Reconstructive surgery should be performed by a urologist, gynecologist, plastic surgeon, or general surgeon who has specialized competence and training in this field.

### Screening

Age-appropriate screening for breast cancer and cervical cancer should be continued unless mastectomy or removal of the cervix has occurred. For patients using androgen therapy who have not had a complete hysterectomy, there may be an increased risk of endometrial cancer and ovarian cancer.

## **Male-to-Female Transgender Individuals**

### Hormones

Estrogen therapy results in gynecomastia, reduced hair growth, redistribution of fat, and reduced testicular volume. All patients considering therapy should be screened for medical contraindications. After surgery, doses of estradiol, 2–4 mg/d, or conjugated equine estrogen, 2.5 mg/d, are often sufficient to keep total testosterone levels to normal female levels of less than 25 ng/dL. Nonoral therapy also can be offered. It is recommended that male-to-female transgender patients receiving estrogen therapy have an annual prolactin level assessment and visual field examination to screen for prolactinoma.

### Surgery

Surgery usually involves penile and testicular excision and the creation of a neovagina. Reported complications of surgery include vaginal and urethral stenosis, fistula formation, problems with remnants of erectile tissue, and pain. Vaginal dilation of the neovagina is required to maintain patency. Other surgical procedures that may be performed include breast implants and nongenital surgery, such as facial feminization surgery.

### Screening

Age-appropriate screening for breast and prostate cancer is appropriate for male-to-female transgender patients. Opinion varies regarding the need for Pap testing in this population. In patients who have a neocervix created from the glans penis, routine cytologic examination of the neocervix may be indicated. The glands are more prone to cancerous changes than the skin of the penile shaft, and intraepithelial neoplasia of the glans is more likely to progress to invasive carcinoma than is intraepithelial neoplasia of other penile skin.

## **SUMMARY**

For member contracts subject to Washington’s Gender Affirming Treatment Act (SSB 5313)

For member contracts subject to the Washington Gender Affirming Treatment Act (SSB 5313), criteria for gender affirming interventions are based on the research, guidelines developed using the available evidence and expert clinical consensus, and on the Act.

---

Therefore, for member contracts subject to the Washington Gender Affirming Treatment Act (SSB 5313), gender affirming interventions for gender dysphoria may be considered medically necessary when specified policy criteria are met.

For member contracts subject to the Washington Gender Affirming Treatment Act (SSB 5313), criteria for gender affirming interventions are based on the Act and on guidelines developed using the available evidence and expert clinical consensus. Therefore, for these members, when these criteria are not met, gender affirming interventions for gender dysphoria are considered not medically necessary.

For member contracts *not* subject to Washington's Gender Affirming Treatment Act (SSB 5313)

The research lacks well-designed studies comparing the safety and effectiveness of no intervention for gender dysphoria with interventions such as gender affirming surgery. However, there are challenges in conducting large studies to evaluate existing treatments, and such studies are not expected in the near future. Although additional research is needed, the research has consistently suggested significant improvement in symptoms and overall quality of life in those who have received certain interventions for gender dysphoria. The World Professional Association for Transgender Health (WPATH) Standards of Care (SOC) for the Health of Transsexual, Transgender, and Gender Nonconforming Peoples recommend that specific criteria are met prior to surgical interventions for gender dysphoria. These guidelines are based on evidence and expert clinical consensus and the included criteria were developed to promote optimal patient care. Therefore, gender affirming interventions for gender dysphoria may be considered medically necessary when specified policy criteria are met.

The World Professional Association for Transgender Health (WPATH) Standards of Care (SOC) for the Health of Transsexual, Transgender, and Gender Nonconforming Peoples recommend that specific criteria are met prior to surgical interventions for gender dysphoria. These guidelines are based on evidence and expert clinical consensus and the included criteria were developed to promote optimal patient care. Therefore, when these criteria are not met, gender affirming interventions for gender dysphoria are considered not medically necessary.

There are no evidence-based clinical practice guidelines that recommend gender affirming surgical interventions not listed in Criterion I.B.2. or revision to a previous gender affirming surgery because of dissatisfaction with the appearance improve health outcomes. Therefore, gender affirming surgical interventions not listed in Criterion I.B.2. and revision to a previous gender affirming surgery because of dissatisfaction with the appearance are considered not medically necessary.

The World Professional Association for Transgender Health (WPATH) Standards of Care (SOC) for the Health of Transsexual, Transgender, and Gender Nonconforming Peoples describe reversible and irreversible interventions, and the ideal order and timing of these approaches. Surgery as an intervention is considered irreversible by WPATH. Therefore, reversal of gender affirming surgery for gender dysphoria is considered not medically necessary.

---

## REFERENCES

1. Position Statement on Access to Care for Transgender and Gender Variant Individuals. Official Position of the American Psychiatric Association. Approved 2018. [cited 11/15/2021]. Available from: <https://www.psychiatry.org/File%20Library/About-APA/Organization-Documents-Policies/Policies/Position-2018-Access-to-Care-for-Transgender-and-Gender-Diverse-Individuals.pdf>.
2. World Professional Association for Transgender Health (WPATH). Position Statement on Medical Necessity of Treatment, Sex Reassignment, and Insurance Coverage in the U.S.A. 21 December 2016. [cited 11/15/2021]. Available from: <https://www.wpath.org/newsroom/medical-necessity-statement>.
3. Anton, Barry S. (2009). Proceedings of the American Psychological Association for the legislative year 2008: Minutes of the annual meeting of the Council of Representatives, February 22-24, 2008, Washington, DC, and August 13 and 17, 2008, Boston, MA, and minutes of the February, June, August, and December 2008 meetings of the Board of Directors. *American Psychologist* 64, 372-453. doi:10.1037/a0015932. PMID:
4. World Professional Association for Transgender Health (WPATH). Standards of Care, Version 7. Standards of Care for the Health of Transsexual, Transgender, and Gender-Nonconforming People. 2011. [cited 11/15/2021]. Available from: [https://s3.amazonaws.com/amo\\_hub\\_content/Association140/files/Standards%20of%20Care%20V7%20-%202011%20WPATH%20\(2\)\(1\).pdf](https://s3.amazonaws.com/amo_hub_content/Association140/files/Standards%20of%20Care%20V7%20-%202011%20WPATH%20(2)(1).pdf).
5. AMAHo Delegates. Resolution #114. American Medical Association House of Delegates. "Removing Barriers to Care for Transgender Patients" [cited. Available from:
6. DHHS. Nondiscrimination in Health Programs and Activities. Department of Health and Human Services. Office of the Secretary. 45 CFR Part 92. RIN 0945-AA02. [cited 2/22/2021]. Available from: <https://s3.amazonaws.com/public-inspection.federalregister.gov/2016-11458.pdf>.
7. American Psychiatric Association (2013): Diagnostic and Statistical Manual of Mental Disorders, 5th ed. Arlington, VA: American Psychiatric Press.
8. S Bucci, G Mazzon, G Liguori, et al. Neovaginal prolapse in male-to-female transsexuals: an 18-year-long experience. *BioMed research international*. 2014;2014:240761. PMID: 24895556
9. G Selvaggi, C Dhejne, M Landen, A Elander. The 2011 WPATH Standards of Care and Penile Reconstruction in Female-to-Male Transsexual Individuals. *Advances in urology*. 2012;2012:581712. PMID: 22654902
10. WC Hembree, PT Cohen-Kettenis, L Gooren, et al. Endocrine Treatment of Gender-Dysphoric/Gender-Incongruent Persons: An Endocrine Society\* Clinical Practice Guideline. *J Clin Endocrinol Metab*. 2017. PMID: 28945902
11. JU Berli, L Capitán, D Simon, R Bluebond-Langner, E Plemons, SD Morrison. Facial gender confirmation surgery—review of the literature and recommendations for Version 8 of the WPATH Standards of Care. *International Journal of Transgenderism*. 2017;18(3):264-70. PMID:
12. MH Murad, MB Elamin, MZ Garcia, et al. Hormonal therapy and sex reassignment: a systematic review and meta-analysis of quality of life and psychosocial outcomes. *Clin Endocrinol (Oxf)*. 2010;72:214-31. PMID: 19473181
13. MB Elamin, MZ Garcia, MH Murad, PJ Erwin, VM Montori. Effect of sex steroid use on cardiovascular risk in transsexual individuals: a systematic review and meta-analyses. *Clin Endocrinol (Oxf)*. 2010;72:1-10. PMID: 19473174

14. AL de Vries, JK McGuire, TD Steensma, EC Wagenaar, TA Doreleijers, PT Cohen-Kettenis. Young adult psychological outcome after puberty suppression and gender reassignment. *Pediatrics*. 2014;134:696-704. PMID: 25201798
15. SD Morrison, F Capitán-Cañadas, A Sánchez-García, et al. Prospective Quality-of-Life Outcomes after Facial Feminization Surgery: An International Multicenter Study. *Plastic and reconstructive surgery*. 2020;145(6):1499-509. PMID: 32459779
16. C Imbimbo, P Verze, A Palmieri, et al. A report from a single institute's 14-year experience in treatment of male-to-female transsexuals. *J Sex Med*. 2009;6:2736-45. PMID: 19619147
17. D Duisin, G Nikolic-Balkoski, B Batinic. Sociodemographic profile of transsexual patients. *Psychiatria Danubina*. 2009;21(2):220-3. PMID: 19556952
18. G Heylens, C Verroken, S De Cock, G T'Sjoen, G De Cuypere. Effects of different steps in gender reassignment therapy on psychopathology: a prospective study of persons with a gender identity disorder. *J Sex Med*. 2014;11(1):119-26. PMID: 24344788
19. AD Fisher, E Bandini, H Casale, et al. Sociodemographic and clinical features of gender identity disorder: an Italian multicentric evaluation. *J Sex Med*. 2013;10(2):408-19. PMID: 23171237
20. A Gorin-Lazard, K Baumstarck, L Boyer, et al. Hormonal therapy is associated with better self-esteem, mood, and quality of life in transsexuals. *J Nerv Ment Dis*. 2013;201:996-1000. PMID: 24177489
21. E Gomez-Gil, L Zubiaurre-Elorza, I Esteva, et al. Hormone-treated transsexuals report less social distress, anxiety and depression. *Psychoneuroendocrinology*. 2012;37:662-70. PMID: 21937168
22. A Johansson, E Sundbom, T Hojerback, O Bodlund. A five-year follow-up study of Swedish adults with gender identity disorder. *Archives of sexual behavior*. 2010;39(6):1429-37. PMID: 19816764
23. H Asscheman, EJ Giltay, JA Megens, WP de Ronde, MA van Trotsenburg, LJ Gooren. A long-term follow-up study of mortality in transsexuals receiving treatment with cross-sex hormones. *Eur J Endocrinol*. 2011;164:635-42. PMID: 21266549
24. Committee Opinion no. 512: health care for transgender individuals. *Obstet Gynecol*. 2011;118:1454-8. PMID: 22105293
25. Committee Opinion No. 685: Care for Transgender Adolescents. *Obstet Gynecol*. 2017;129:e11-e16. PMID: 28002311

## CODES

### NOTES:

- Follicular unit extraction (FEU) of individual hairs is correctly coded with code 15775 or 15776 and is determined by the number of "punch grafts" performed. Be advised that standard CMS Medically Unlikely Edits (MUEs or Maximum Units of Service) will apply.
- Code 17999 should be reported for laser hair removal. This code may also be used for abdominoplasty or calf/pectoral implants.
- Codes 31552, 31554, 31580, 31584, 31587, and 31591 are not appropriate to use to represent voice modification. Unlisted code 31599 should be reported instead.
- Code 31899 should be reported for reduction thyroid chondroplasty (reduction of the thyroid cartilage or Adam's Apple).
- Code 40799 should be reported for lip reduction.
- Code 55899 should be reported for phallic reconstruction/phalloplasty.

- Codes 55970 and 55980 are non-specific. The specific procedure code(s) must be requested in place of these non-specific codes.

<b>Codes</b>	<b>Number</b>	<b>Description</b>
CPT	11950	Subcutaneous injection of filling material (eg, collagen); 1 cc or less
	11951	Subcutaneous injection of filling material (eg, collagen); 1.1 to 5.0 cc
	11952	Subcutaneous injection of filling material (eg, collagen); 5.1 to 10.0 cc
	11954	Subcutaneous injection of filling material (eg, collagen); over 10 cc
	11970	Replacement of tissue expander with permanent implant
	11971	Removal of tissue expander(s) without insertion of implant
	15775	Punch graft for hair transplant; 1 to 15 punch grafts
	15776	Punch graft for hair transplant; more than 15 punch grafts
	15820	Blepharoplasty, lower eyelid
	15821	;with extensive herniated fat pad
	15822	Blepharoplasty, upper eyelid
	15823	;with excessive skin weighting down lid
	15824	Rhytidectomy; forehead
	15825	Rhytidectomy; neck with platysmal tightening (platysmal flap, P-flap)
	15826	Rhytidectomy; glabellar frown lines
	15828	Rhytidectomy; cheek, chin, and neck
	15829	Rhytidectomy; superficial musculoaponeurotic system (SMAS) flap
	15830	Excision, excessive skin and subcutaneous tissue (includes lipectomy); abdomen, infraumbilical panniculectomy
	15832	Excision, excessive skin and subcutaneous tissue (includes lipectomy); thigh
	15833	Excision, excessive skin and subcutaneous tissue (includes lipectomy); leg
	15834	Excision, excessive skin and subcutaneous tissue (includes lipectomy); hip
	15835	Excision, excessive skin and subcutaneous tissue (includes lipectomy); buttock
	15836	Excision, excessive skin and subcutaneous tissue (includes lipectomy); arm
	15837	Excision, excessive skin and subcutaneous tissue (includes lipectomy); forearm or hand
	15838	Excision, excessive skin and subcutaneous tissue (includes lipectomy); submental fat pad
	15839	Excision, excessive skin and subcutaneous tissue (includes lipectomy); other area
	15847	Excision, excessive skin and subcutaneous tissue (includes lipectomy), abdomen (eg, abdominoplasty) (includes umbilical transposition and fascial plication) (List separately in addition to code for primary procedure)
	15876	Suction assisted lipectomy; head and neck
	15877	Suction assisted lipectomy; trunk
	15878	Suction assisted lipectomy; upper extremity
	15879	Suction assisted lipectomy; lower extremity
	17380	Electrolysis epilation, each 30 minutes
	17999	Unlisted procedure, skin, mucous membrane and subcutaneous tissue
	19303	Mastectomy, simple, complete
	19316	Mastopexy
	19318	Breast reduction
	19325	Breast augmentation with implant
	19350	Nipple/areola reconstruction
	19499	Unlisted procedure, breast
	21120	Genioplasty; augmentation (autograft, allograft, prosthetic material)
	21121	Genioplasty; sliding osteotomy, single piece
	21122	Genioplasty; sliding osteotomies, 2 or more osteotomies (eg, wedge excision or bone wedge reversal for asymmetrical chin)

21123	Genioplasty; sliding, augmentation with interpositional bone grafts (includes obtaining autografts)
21209	Osteoplasty, facial bones; reduction
21270	Malar augmentation, prosthetic material
30400	Rhinoplasty, primary; lateral and alar cartilages and/or elevation of nasal tip
30410	;complete, external parts including bony pyramid, lateral and alar cartilages, and/or elevation of nasal tip
30420	;including major septal repair
30430	Rhinoplasty, secondary; minor revision (small amount of nasal tip work)
30435	;intermediate revision (bony work with osteotomies)
30450	;major revision (nasal tip work and osteotomies)
31599	Unlisted procedure, larynx
31899	Unlisted procedure, trachea, bronchi
40799	Unlisted procedure, lips
53400	Urethroplasty; first stage, for fistula, diverticulum, or stricture (eg, Johanssen type)
53405	Urethroplasty; second stage (formation of urethra), including urinary diversion
53410	Urethroplasty, 1-stage reconstruction of male anterior urethra
53415	Urethroplasty, transpubic or perineal, 1-stage, for reconstruction or repair of prostatic or membranous urethra
53420	Urethroplasty, 2-stage reconstruction or repair of prostatic or membranous urethra; first stage
53425	Urethroplasty, 2-stage reconstruction or repair of prostatic or membranous urethra; second stage
53430	Urethroplasty, reconstruction of female urethra
54125	Amputation of penis; complete (Penectomy)
54400	Insertion of penile prosthesis; non-inflatable (semi-rigid)
54401	Insertion of penile prosthesis; inflatable (self-contained)
54405	Insertion of multi-component, inflatable penile prosthesis, including placement of pump, cylinders, and reservoir
54520	Orchiectomy, simple (including subcapsular), with or without testicular prosthesis, scrotal or inguinal approach
54660	Insertion of testicular prosthesis
54690	Laparoscopy, surgical; orchiectomy
55175	Scrotoplasty; simple
55180	Scrotoplasty; complicated
55899	Phallic reconstruction/Phalloplasty (Unlisted procedure, male genital system)
55970	intersex surgery; male to female
55980	intersex surgery; female to male
56625	Vulvectomy simple; complete
56800	Plastic repair of introitus
56805	Clitoroplasty for intersex state
57106	Vaginectomy, partial removal of vaginal wall
57110	Vaginectomy, complete removal of vaginal wall;
57291	Construction of artificial vagina; without graft
57292	Construction of artificial vagina; with graft
57295	Revision (including removal) of prosthetic vaginal graft; vaginal approach
57296	Revision (including removal) of prosthetic vaginal graft; open abdominal approach
57335	Vaginoplasty for intersex state
57426	Revision (including removal) of prosthetic vaginal graft, laparoscopic approach

58150	Total abdominal hysterectomy (corpus and cervix), with or without removal of tube(s), with or without removal of ovary(s)
58180	Supracervical abdominal hysterectomy (subtotal hysterectomy), with or without removal of tube(s), with or without removal of ovary(s)
58260	Vaginal hysterectomy, for uterus 250 g or less
58262	Vaginal hysterectomy, for uterus 250 g or less; with removal of tube(s), and/or ovary(s)
58270	Vaginal hysterectomy, for uterus 250 g or less; with repair of enterocele
58275	Vaginal hysterectomy, with total or partial vaginectomy;
58290	Vaginal hysterectomy, for uterus greater than 250 g
58291	Vaginal hysterectomy, for uterus greater than 250 g; with removal of tube(s) and/or ovary(s)
58353	Endometrial ablation, thermal, without hysteroscopic guidance
58356	Endometrial cryoablation with ultrasonic guidance, including endometrial curettage, when performed
58541	Laparoscopy, surgical, supracervical hysterectomy, for uterus 250 g or less
58542	Laparoscopy, surgical, supracervical hysterectomy, for uterus 250 g or less; with removal of tube(s) and/or ovary(s)
58543	Laparoscopy, surgical, supracervical hysterectomy, for uterus greater than 250 g
58544	Laparoscopy, surgical, supracervical hysterectomy, for uterus greater than 250 g; with removal of tube(s) and/or ovary(s)
58550	Laparoscopy, surgical, with vaginal hysterectomy, for uterus 250 g or less
58552	Laparoscopy, surgical, with vaginal hysterectomy, for uterus 250 g or less; with removal of tube(s) and/or ovary(s)
58553	Laparoscopy, surgical, with vaginal hysterectomy, for uterus greater than 250 g
58554	Laparoscopy, surgical, with vaginal hysterectomy, for uterus greater than 250 g; with removal of tube(s) and/or ovary(s)
58563	Hysteroscopy, surgical; with endometrial ablation (eg. Endometrial resection, electro-surgical ablation, thermoablation)
58570	Laparoscopy, surgical, with total hysterectomy, for uterus 250 g or less
58571	Laparoscopy, surgical, with total hysterectomy, for uterus 250 g or less; with removal of tube(s) and/or ovary(s)
58572	Laparoscopy, surgical, with total hysterectomy, for uterus greater than 250 g
58573	Laparoscopy, surgical, with total hysterectomy, for uterus greater than 250 g; with removal of tube(s) and/or ovary(s)
58720	Salpingo-oophorectomy, complete or partial, unilateral or bilateral (separate procedure)
67900	Repair of brow ptosis (supraciliary, mid-forehead or coronal approach)
67901	Repair of blepharoptosis; frontalis muscle technique with suture or other material (eg, banked fascia)
67902	Repair of blepharoptosis; frontalis muscle technique with autologous fascial sling (includes obtaining fascia)
67903	Repair of blepharoptosis; (tarso) levator resection or advancement, internal approach
67904	Repair of blepharoptosis; (tarso) levator resection or advancement, external approach
67906	Repair of blepharoptosis; superior rectus technique with fascial sling (includes obtaining fascia)
67908	Repair of blepharoptosis; conjunctivo-tarso-Muller's muscle-levator resection (eg, Fasanella-Servat type)
67909	Reduction of overcorrection of ptosis

	67950	Canthoplasty (reconstruction of canthus)
HCPCS	C1813	Prosthesis, penile, inflatable
	C2622	Prosthesis, penile, noninflatable
	L8039	Breast prosthesis, not otherwise specified
	L8600	Implantable breast prosthesis, silicone or equal

**Date of Origin:** September 2014